A constructive proof of the Wedderburn-Artin theorem

Sheng Gao * †

Department of Mathematics, Hefei University of Technology, Hefei 230026, China

Abstract. In this short note, we use the idempotent decomposition to give an explicit isomorphism from a semisimple Artinian ring to an external direct sum of finite full matrix rings over division rings. In particular, we directly construct the matrix units in a semisimple Artinian ring.
Key Words: semisimple Artinian rings; complete reducible modules; idempotents; right ideals.
2000 Mathematics Subject Classification: 16P20.

arXiv:1403.4731v1 [math.RT] 19 Mar 2014

Throughout this note A denotes a ring with identity. If e and f are two idempotents of A such that eA and fA are isomorphic as right A-module, then e and f are said to be equivalent and we write $e \simeq f$. Concerning the idempotents, all the facts needed for our discussion are collected in the following lemma.

Lemma 1. [6, Theorem 4.1-4.4, pp.16-18] *Let e and f be idempotents of A.*

(1) *If $(eA)_A$ is an internal direct sum of A-submodules $eA = I_1 \oplus I_2 \oplus \cdots \oplus I_n$, then there exists an orthogonal idempotent decomposition $e = e_1 + e_2 + \cdots + e_n$ such that $I_i = e_iA$ $(1 \leq i \leq n)$. e is primitive if and only if $(eA)_A$ is indecomposable. In particular, if $(eA)_A$ is irreducible, then e is primitive.*

(2) *$e \simeq f$ if and only if there exist $a \in fAe, b \in eAf$ such that $ab = f, ba = e$.*

(3) *$\mathrm{Hom}_A(eA, fA) \cong fAe$ as additive groups and $\mathrm{End}_A(eA) \cong eAe$ as rings.*

Lemma 2. [3, Lemma 1] *If A contains a family elements $\varepsilon_{\mu\nu}(1 \leq \mu, \nu \leq n)$ such that $\varepsilon_{11} + \cdots + \varepsilon_{nn} = 1_A$ and $\varepsilon_{\mu\nu}\varepsilon_{\xi\eta} = \delta_{\nu\xi}\varepsilon_{\mu\eta}$ for all $\mu, \nu, \xi, \eta \in \{1, \cdots, n\}$(here $\delta_{\nu\xi}$ is the Kronecker symbol), then we have the following ring isomorphism*

$$\begin{aligned} \varphi : A &\longrightarrow M_n(\varepsilon_{11}A\varepsilon_{11}), \\ a &\longmapsto (\varepsilon_{1\mu}a\varepsilon_{\nu 1})_{n\times n} \end{aligned}$$

where the (μ, ν)-entry of $\varphi(a)$ is $\varepsilon_{1\mu}a\varepsilon_{\nu 1}$. (The elements $\varepsilon_{\mu\nu}$'s are called matrix units in A.)

Now, we will prove the main result of this article.

Wedderburn-Artin's Theorem. *If the regular right A-module A_A is completely reducible, then A is isomorphic to an external direct sum of finite full matrix rings over division rings.*

Proof. We can express A_A as an internal direct sum of a family of irreducible A-submodules: $A = \bigoplus_{\lambda\in\Lambda} I_\lambda$. Then there exist finite elements $\lambda_1, \lambda_2, \cdots, \lambda_n \in \Lambda$ such that $1_A \in \bigoplus_{i=1}^n I_{\lambda_i}$. This yields that $A = \bigoplus_{i=1}^n I_{\lambda_i}$ and $\Lambda = \{\lambda_1, \lambda_2, \ldots, \lambda_n\}$. Thus 1_A has a orthogonal primitive idempotent decomposition $1_A = \sum_{i=1}^k \sum_{\mu=1}^{n_i} e_{i,\mu}$, where $e_{i,\mu}$'s are pairwise orthogonal primitive idempotents of A such that $e_{i,\mu} \simeq e_{j,\nu}$

*E-mail address: highlyrise@aliyun.com
†This manuscript has been submitted to the Journal of University of Science and Technology of China.

holds if and only if $i = j$. Furthermore, for each i, μ with $1 \leq i \leq k$ and $1 \leq \mu \leq n_i$, we choose $a_\mu^{(i)} \in e_{i,1}Ae_{i,\mu}$ and $b_\mu^{(i)} \in e_{i,\mu}Ae_{i,1}$ such that $e_{i,1} = a_\mu^{(i)} b_\mu^{(i)}$ and $e_{i,\mu} = b_\mu^{(i)} a_\mu^{(i)}$. In particular, set $a_1^{(i)} = b_1^{(i)} = e_{i,1}$.By Schur's lemma, $e_{i,1}Ae_{i,1} \cong \mathrm{End}_A(e_{i,1}A)$ is a division ring for each i and $e_{i,\mu}Ae_{j,\nu} \cong \mathrm{Hom}_A(e_{j,\nu}A, e_{i,\mu}A) = 0$ whenever $i \neq j$. For every $i (1 \leq i \leq k)$, set $c_i = \sum_{\mu=1}^{n_i} e_{i,\mu}$. Then $1_A = \sum_{i=1}^{k} c_i$ and $c_i A c_j = 0$ whenever $i \neq j$, which yields that $c_i^2 = c_i \sum_{j=1}^{k} c_j = c_i$ and $c_i a = c_i a \sum_{j=1}^{k} c_j = c_i a c_i = \sum_{j=1}^{k} c_j a c_i = a c_i$ for every $a \in A$. Hence c_i's are all central idempotents of A and they are pairwise orthogonal. From this fact we can conclude that $A = \bigoplus_{i=1}^{k} Ac_i$, where each Ac_i is a two-sided ideal of A and is itself a ring with identity c_i. Set $A_i = Ac_i = c_i A$.

Using Lemma 2, we can easily construct an isomorphism from A_i to the $n_i \times n_i$ full matrix ring over $e_{i,1}Ae_{i,1}$. For each $\mu \in \{1, \cdots, n_i\}$, we have $e_{i,\mu} = e_{i,\mu} \sum_{\nu=1}^{n_i} e_{i,\nu} = e_{i,\mu}c_i = c_i e_{i,\mu} \in A_i$, so $\{b_\mu^{(i)}, a_\nu^{(i)} | 1 \leq \mu, \nu \leq n_i\} \subset A_i$. An easy computation shows that $\sum_{\mu=1}^{n_i} b_\mu^{(i)} a_\mu^{(i)} = \sum_{\mu=1}^{n_i} e_{i,\mu} = c_i = 1_{A_i}$ and $(b_\mu^{(i)} a_\nu^{(i)})(b_\xi^{(i)} a_\eta^{(i)}) = \delta_{\nu\xi} b_\mu^{(i)} a_\eta^{(i)}$ for $\mu, \nu, \eta, \xi \in \{1, \cdots, n_i\}$. Thus $b_\mu^{(i)} a_\nu^{(i)} (1 \leq \mu, \nu \leq n_i)$ are matrix units in A_i, which yields the following isomorphism of rings:

$$\begin{aligned} \Upsilon_i : A_i &\longrightarrow M_{n_i}(e_{i,1}Ae_{i,1}), \\ ac_i &\longmapsto \left(b_1^{(i)} a_\mu^{(i)} a c_i b_\nu^{(i)} a_1^{(i)}\right)_{n_i \times n_i} = \left(a_\mu^{(i)} a b_\nu^{(i)}\right)_{n_i \times n_i} \end{aligned}$$

where a is any element of A and $\left(a_\mu^{(i)} a b_\nu^{(i)}\right)_{n_i \times n_i}$ denotes a $n_i \times n_i$-matrix whose (μ, ν)-entry is $a_\mu^{(i)} a b_\nu^{(i)}$ $(1 \leq \mu, \nu \leq n_i)$. As a result, we obtain the following isomorphism of rings:

$$\begin{aligned} \Upsilon : A &\longrightarrow M_{n_1}(e_{1,1}Ae_{1,1}) \bigoplus \cdots \bigoplus M_{n_k}(e_{k,1}Ae_{k,1}) \ . \\ a &\longmapsto \left(\begin{pmatrix} a_1^{(1)} a b_1^{(1)}, & \cdots, & a_1^{(1)} a b_{n_1}^{(1)} \\ \cdots & \cdots & \cdots \\ a_{n_1}^{(1)} a b_1^{(1)}, & \cdots, & a_{n_1}^{(1)} a b_{n_1}^{(1)} \end{pmatrix}, \cdots, \begin{pmatrix} a_1^{(k)} a b_1^{(k)}, & \cdots, & a_1^{(k)} a b_{n_k}^{(k)} \\ \cdots & \cdots & \cdots \\ a_{n_k}^{(k)} a b_1^{(k)}, & \cdots, & a_{n_k}^{(k)} a b_{n_k}^{(k)} \end{pmatrix} \right) \end{aligned}$$

□

ACKNOWLEDGMENTS

I would like to express my gratitude to Prof.J.P.Zhang, who led me to the fascinating field of representation theory. Thanks are also given to Prof.Nicholson for his helpful comments and suggestions.

References

[1] E.Artin, C.J.Nesbitt and R.M.Thrall, rings with minimum condition, university of Michigan Press, 1946.
[2] E.Artin and G.Whaples, The theory of simple rings, American J.Math.65(1943), No.1, 87-107.
[3] M.Bresar, An elementary approach to Wedderburn's structure theory, Expo.Math.28(2010), 79-83.
[4] N.Jacobson, Basic Algebra II, W.H.Freeman and Company, 1989.
[5] Shaoxue Liu, Wedderburn-Artin theorem for additive categories, Kexue Tongbao(English Ed.)33(1988), No.18, 1501-1503.
[6] H.Nagao and Y.Tsushima, Representations of Finite Groups, Academic Press, New York, 1987.
[7] W.K.Nicholson, A short proof of the Wedderburn-Artin theorem, New Zealand J.Math.22(1993), No.1, 83-86.
[8] T.Szele, Simple proof of the Wedderburn-Artin structure theorem, Acta Math.Acad.Sci.Hungar.5(1954), 101-107.

[9] Bangteng Xu, A simplified and elementary proof of the Wedderburn-Artin structure theorem(in Chinese), Hunan Jiaoyu Xueyuan Xuebao(Ziran Kexue)8(1990), No.5, 143-145.